

### Self-similar solution of the mixing layer

The starting point for this exercise is the formal expression of the self-similar solution for the plane mixing layer flow, established in the course slides (you don't have to demonstrate all those results again).

1. Show that for a free plane mixing layer

$$\bar{U}_2 = \Delta U \delta' \left\{ \eta f - \int_0^\eta f d\tilde{\eta} \right\}$$

2. From the simplified equation governing the mean flow, show that the self-similarity imposes a linear growth of the mixing layer, denoted  $\delta = a(x_1 - x_1^0)$  with  $a$  and  $x_1^0$  two constants ( $x_1^0$  represents the virtual spatial origin of the self-similar flow).
3. Show that  $S\eta f' = g'$  where  $S$  is the spreading rate of the mixing layer

$$S \equiv \frac{u_m}{\Delta U} \delta'$$

4. The Reynolds stress tensor is modeled by introducing a turbulent viscosity  $\nu_t$  (Boussinesq model), and it will be assumed that  $\nu_t = l_m \Delta U$  where  $l_m = \alpha \delta(x_1)$  is a mixing length scale. Justify in a few sentences this expression of the turbulent viscosity.
5. Show that the function  $f$  must satisfied the following differential equation,  $f'' = -S \text{Re}_t \eta f'$ , where  $\text{Re}_t$  denotes a turbulent Reynolds number.
6. By integrating the previous equation, in taking into account the associated boundary conditions, show finally that

$$f(\eta) = \frac{1}{2} \text{erf}\left(\frac{\eta}{\sqrt{2}\sigma}\right) \quad \text{where} \quad \text{erf}(\xi) = \frac{2}{\sqrt{\pi}} \int_0^\xi e^{-\zeta^2} d\zeta$$

is the error function.

7. Determine an estimate of the values of  $S$  and  $a$  the measurements by Champagne *et al.*<sup>2</sup> and by Bell & Mehta<sup>1</sup> (these papers can be downloaded from our website).

### To be finish as homework

8. Plot the curve  $(\bar{U}_1 - u_m)/\Delta U$  as a function of  $\eta$  and compare to experimental data.
9. Compare also the self-similar solution with the tanh profile used to solve Rayleigh's equation in a previous homework. Include this plot in your copy (homework on Stability).

### References

- <sup>1</sup> Bell, J.H. & Mehta, J.M., 1990, Development of a two-stream mixing layer from tripped and untripped boundary layers, *AIAA Journal*, **28**(12), 2034-2042.
- <sup>2</sup> Champagne, F.H., Pao, Y.H. & Wygnanski, I.J., 1976, On the two-dimensional mixing region *J. Fluid Mech.*, **74**(2), 209-250.
- <sup>3</sup> Pope, S.B., 2000, *Turbulent flows*, Cambridge University Press.